




Angles and Degree Measure

An **angle** can be formed by rotating one ray away from a fixed ray indicated by an arrow. The fixed ray is the **initial side** and the rotated ray is the **terminal side**. An angle whose vertex is the center of a circle is a **central angle**, and the arc of the circle through which the terminal side moves is the **intercepted arc**. An angle in **standard position** is located in a rectangular coordinate system with the vertex at the origin and the initial side on the positive x-axis.

Angle α 

Central angle



Angle in standard position

Degree Measure of Angles



The measure, $m(\alpha)$, of an angle α is the amount of rotation from the initial side to the terminal side, and is found by using any circle centered at the vertex. An angle that forms a complete circle arc is 360° .

The **degree measure of an angle** is the number of degrees in the intercepted arc of a circle centered at the vertex.


Counterclockwise rotation—positive angle

Clockwise rotation—negative angle


An angle in standard position is said to lie in the quadrant where its terminal side lies.




Acute angle




Obtuse angle



Right angle



Straight angle



Quadrantal angle

Acute angle—An angle with a measure between 0° and 90° .

Obtuse angle—An angle with a measure between 90° and 180° .

Straight angle—An angle with a measure of exactly 180° .

Right angle—An angle with a measure of exactly 90° .

Quadrantal angle—An angle in standard position whose terminal side is on an axis.

The terminal side of an angle may be rotated in either a positive or negative direction to get to its final position. It can also be rotated for more than one revolution in either direction. Two angles with the same terminal side are called **coterminal angles**.

Coterminal Angles—Angles α and β are coterminal if and only if there is an integer k such that

$$m(\beta) = m(\alpha) + k360^\circ$$

Add or subtract multiples of 360°

Examples: Find two positive angles and two negative angles that are coterminal with each angle:

a) 23°

$$23^\circ - 360^\circ = -337^\circ$$

$$23^\circ + 360^\circ = 383^\circ$$

$$-337^\circ - 360^\circ = -697^\circ$$

$$383^\circ + 360^\circ = 743^\circ$$

b) -146°

$$-146^\circ - 360^\circ = -506^\circ$$

$$-146^\circ + 360^\circ = 214^\circ$$

$$-506^\circ - 360^\circ = -866^\circ$$

$$214^\circ + 360^\circ = 574^\circ$$

Examples: Determine whether the angles in each pair are coterminal:

a) -128° and 592°

$$592^\circ - (-128^\circ) = 720^\circ$$

720° is a multiple of 360° . **Coterminal**

b) 8° and -368°

$$8 - (-368) = 376^\circ$$

376° is not a multiple of 360° . **Not coterminal**

Example: Name the quadrant in which each angle lies.

a) 255°



b) -650°

$$-650^\circ + 360^\circ = -290^\circ$$

$$-290^\circ + 360^\circ = 70^\circ$$

c) 1360°

$$1360^\circ - 360^\circ = 1000^\circ$$

$$1000^\circ - 360^\circ = 640^\circ$$

$$640^\circ - 360^\circ = 280^\circ$$

Minutes and Seconds

Historically, angles were measured by using the **degrees-minutes-seconds (DMS) format**, but with calculators it is convenient to have some fractional parts of degrees written in decimal form. Each degree is divided into 60 equal parts called **minutes** (n'), and each minute is divided into 60 equal parts called **seconds** (n'').

$$1 \text{ degree} = 60 \text{ minutes}$$

$$1 \text{ minute} = 60 \text{ seconds}$$

$$1 \text{ degree} = 3600 \text{ seconds}$$

Examples: Perform the indicated operations.

a) $15^\circ 56' 45'' + 18^\circ 12' 33''$

$$\begin{array}{r} 15^\circ 56' 45'' \\ + 18^\circ 12' 33'' \\ \hline 33^\circ 68' 78'' \text{ change } 60'' \text{ to } 1' \\ 33^\circ 69' 18'' \text{ change } 60' \text{ to } 1^\circ \\ \hline 34^\circ 9' 18'' \end{array}$$

b) $36^\circ 5' - 22^\circ 33' 12''$


$$\begin{array}{r} 36^\circ 5' = 36^\circ 4' 60'' \\ = 35^\circ 64' 60'' \\ - 22^\circ 33' 12'' \\ \hline 13^\circ 31' 48'' \end{array}$$

c) $\frac{15^\circ 56' 45''}{2}$

$$\begin{array}{r} 15^\circ 56' 45'' \\ \div 2 \\ \hline 7.5^\circ 28' 22.5'' \\ .5^\circ = 30' \\ \hline 7^\circ 58' 22.5'' \end{array}$$

Examples: Find the degree measure of angle α in each figure.

a)




$$90^\circ + 48^\circ 9' 6'' = 138^\circ 9' 6''$$

$$180^\circ - 138^\circ 9' 6'' = 41^\circ 50' 54''$$

$\alpha = 41^\circ 50' 54''$

b)



$$90^\circ - 75^\circ 5' 6'' = 14^\circ 54' 54''$$

$\alpha = 14^\circ 54' 54''$

Radian Measure, Arc Length, and Area

Unit circle: A circle of radius one that is centered at the origin.

What is the circumference of the unit circle? (Circumference of a circle: $C = \pi d = 2\pi r$) $r = 1$
 $C = 2\pi(1) = 2\pi$

What is the arc length intercepted by a 180° angle ($1/2$ of the circle)? $(\frac{1}{2})(2\pi) = \pi$

What is the arc length intercepted by a 120° angle ($1/3$ of the circle)? $(\frac{1}{3})(2\pi) = \frac{2\pi}{3}$

What is the arc length intercepted by a 30° angle? An angle of 225° ? An angle of 210° ?


$$\left(\frac{30^\circ}{360^\circ}\right)(2\pi) = \frac{\pi}{6}$$

$$\left(\frac{225^\circ}{360^\circ}\right)(2\pi) = \frac{5\pi}{4}$$



$$\left(\frac{210^\circ}{360^\circ}\right)(2\pi) = \frac{7\pi}{6}$$

You have just calculated the radian measure of each of these angles.

The **radian measure** of the angle α in standard position is the directed length of the intercepted arc on the unit circle. **Directed length** means that the radian measure is positive if the rotation of the terminal side is counterclockwise and negative if the rotation is clockwise.



One radian: The angle that intercepts an arc with length equal to the radius of a circle. (On the unit circle, one radian is the angle that intercepts an arc of length one.)



Since the radius of the unit circle is the real number 1, without any dimension (ft., meters, etc.), the arc length, and hence the radian measure of an angle, is also a real number without any dimension. Thus, one radian (1 rad) is the real number 1.

Converting Between Radians and Degrees:

Since there are 2π radians in a circle (the circumference of the unit circle is 2π) and 360° in a circle,
 2π radians = 360° , or π radians = 180° .

Degrees \rightarrow Radians: multiply by $\frac{\pi \text{ rad}}{180^\circ}$.

Radians \rightarrow Degrees: multiply by $\frac{180^\circ}{\pi \text{ rad}}$.

Examples:

Convert the degree measures to radians:

a) 210°

b) -27.2°

$$\left(\frac{210^\circ}{180^\circ}\right)(\pi \text{ rad}) = \frac{7\pi}{6}$$

$$\left(\frac{-27.2^\circ}{180^\circ}\right)(\pi \text{ rad}) = -.475$$

Convert the radian measures to degrees:

a) $\frac{5\pi}{3}$

b) 16.7 radians

$$\left(\frac{5\pi}{3}\right)\left(\frac{180^\circ}{\pi}\right) = 300^\circ$$

$$(16.7)\left(\frac{180^\circ}{\pi}\right) = 956.8^\circ$$

Arc Length

Often, we want to find the arc length (s) on a circle of radius r , intercepted by an angle α :

Since a central angle of 2π (360°) intercepts an arc whose length is the circumference of a circle, we have:

$$\frac{\text{arc length}}{\text{circumference}} = \frac{s}{2\pi r} = \frac{\alpha \text{ in radians}}{2\pi \text{ radians}} = \frac{\alpha \text{ in degrees}}{360^\circ}$$

If α is in radians, we can multiply both sides of the proportion by $2\pi r$ to obtain the arc-length formula:

The length s of an arc intercepted by a central angle of α radians on a circle of radius r is given by: $s = \alpha r$.

This formula only works if α is in radians!

You can solve arc length problems by using the arc-length formula, proportions, or by noting that arc length is a fraction of the circumference of a circle.

Examples:

A central angle of $\pi/2$ intercepts an arc on the surface of the earth that runs from the equator to the North Pole.

Using 3950 miles as the radius of the earth, find the length of the intercepted arc to the nearest mile.

$$\begin{aligned}\alpha &= \pi/2 \\ r &= 3950 \text{ mi} \\ s &= \alpha r \\ s &= (\pi/2)(3950) = \boxed{6204.6 \text{ mi}}\end{aligned}$$

A wagon wheel has a diameter of 28 inches and an angle of 30° between the spokes. What is the length of the arc s (to the nearest hundredth of an inch) between two adjacent spokes?

$$\begin{aligned}\alpha &= 30^\circ = \pi/6 \\ r &= 14 \text{ in} \\ s &= \alpha r \\ s &= (\pi/6)(14) = \boxed{7.3 \text{ in}}\end{aligned}$$

Area of a Sector of a Circle

Finding the area (A) of a sector of a circle of radius r with central angle α , is similar to finding the arc length:

$$\frac{\text{area of sector}}{\text{area of circle}} = \frac{A}{\pi r^2} = \frac{\alpha \text{ in radians}}{2\pi \text{ radians}} = \frac{\alpha \text{ in degrees}}{360^\circ}$$

If α is in radians, we can multiply both sides of the proportion by πr^2 to obtain a formula for the area of the

sector of a circle: $A = \frac{\alpha r^2}{2}$. **Once again, this formula only works if α is in radians!**

Examples:

Which is bigger: a slice of pizza from a 10" diameter pizza cut into 6 slices, or a slice from a 12" diameter pizza cut into 8 slices?

$$\begin{aligned}\text{Area of 10" diameter pizza: } A &= \pi(5)^2 = 25\pi \\ \text{Area of } 1/6 \text{ of pizza: } \frac{25\pi}{6} &= 13.1 \text{ in}^2 \\ \text{Area of 12" diameter pizza: } A &= \pi(6)^2 = 36\pi \\ \text{Area of } 1/8 \text{ of pizza: } \frac{36\pi}{8} &= 14.1 \text{ in}^2\end{aligned}$$

A center-pivot irrigation system is used to water a circular field with radius 200 feet. In three hours the system waters a sector with a central angle of $\pi/8$. What area (in square feet) is watered in that time?

$$\begin{aligned}\text{Area of field: } A &= \pi(200)^2 = 125,663.7 \text{ ft}^2 \\ \text{fraction of circle watered} \rightarrow \left(\frac{\pi/8}{2\pi}\right) & \times \text{area of circle} = \boxed{7854.0 \text{ ft}^2} \quad \text{--OR--} \\ A &= \frac{\alpha r^2}{2} = \frac{(\pi/8)(200)^2}{2} = \boxed{7854.0 \text{ ft}^2}\end{aligned}$$

Angular and Linear Velocity

Velocity: The rate at which the location of an object is changing with respect to time.

Angular Velocity: The rate at which the angle is changing. If a point is in motion on a circle through an angle of α radians in time t , then its angular velocity ω is given by $\omega = \frac{\alpha}{t}$. Angular velocity is usually expressed as radians per unit of time (radians/hr, radians/min, radians/sec, etc.)

Examples:

Convert 650 rpm (revolutions per minute) to radians per minute. (Use the fact that 1 revolution = 2π radians).

$$\frac{650 \text{ rev}}{\text{min}} \cdot \frac{2\pi \text{ rad}}{1 \text{ rev}} = \boxed{4,084.1 \text{ rad/min}}$$

Convert the angular velocity of 1600 rad/hr to rad/sec.

$$\frac{1600 \text{ rad}}{\text{hr}} \cdot \frac{\text{hr}}{60 \text{ min}} \cdot \frac{\text{min}}{60 \text{ sec}} = \boxed{.444 \text{ rad/sec}}$$

A 24-inch lawnmower blade rotates at a rate of 2000 rpm. What is the angular velocity in radians per second of a point on the tip of the blade?

$$\frac{2000 \text{ rev}}{\text{min}} \cdot \frac{2\pi \text{ rad}}{1 \text{ rev}} \cdot \frac{\text{min}}{60 \text{ sec}} = \boxed{209.4 \text{ rad/sec}}$$

Find the angular velocity in radians per second for a particle that is moving in a circular path at 4 revolutions per second on a circle of radius 9 ft.

$$\frac{4 \text{ rev}}{\text{sec}} \cdot \frac{2\pi \text{ rad}}{1 \text{ rev}} = \boxed{25.1 \text{ rad/sec}}$$

Linear Velocity: The rate at which the distance is changing. If a point is in motion on a circle of radius r through an angle of α radians in time t , then its linear velocity v is given by $v = \frac{s}{t}$, where s is the arc length determined by $s = ar$.

Examples:

A propeller with a radius of 1.6 meters is rotating at 1500 revolutions per minute. What is the linear velocity in meters per minute for a point on the tip of the propeller?

$$\begin{aligned} r &= 1.6 \text{ m} \\ C &= 2\pi(1.6) = 10.053 \text{ m} \\ &\text{This is distance point travels each revolution} \end{aligned} \quad \frac{1500 \text{ rev}}{\text{min}} \cdot \frac{10.053 \text{ m}}{1 \text{ rev}} = \boxed{15,079.6 \text{ m/min}}$$

Find the linear velocity in meters per second for a particle that is moving in a circular path at 7 revolutions per second on a circle of radius 15 meters.

$$\frac{7 \text{ rev}}{\text{sec}} \cdot \frac{2\pi(15) \text{ m}}{1 \text{ rev}} = \boxed{659.7 \text{ m/sec}}$$

What is the linear velocity in miles per hour of the tip of a 20-inch ^{diameter} lawnmower blade that is rotating at 3000 rpm?
 $C = 20\pi$ in

$$\frac{3000 \text{ rev}}{\text{min}} \cdot \frac{20\pi \text{ in}}{\text{rev}} \cdot \frac{60 \text{ min}}{\text{hr}} \cdot \frac{\text{ft}}{12 \text{ in}} \cdot \frac{\text{mi}}{5280 \text{ ft}} = \boxed{178.5 \text{ mph}}$$

Find the linear velocity in miles per hour for a particle that is moving in a circular path at 1800 revolutions per minute on a circle with a diameter of 14 inches. $C = 14\pi$ in

$$\frac{1800 \text{ rev}}{\text{min}} \cdot \frac{14\pi \text{ in}}{\text{rev}} \cdot \frac{60 \text{ min}}{\text{hr}} \cdot \frac{\text{ft}}{12 \text{ in}} \cdot \frac{\text{mi}}{5280 \text{ ft}} = \boxed{75.0 \text{ mph}}$$

Linear Velocity in Terms of Angular Velocity: If v is the linear velocity of a point on a circle of radius r , and ω is its angular velocity, then $v = r\omega$.


Example:

Any point on the surface of the earth (except at the poles) makes one revolution (2π radians) about the axis of the earth in 24 hours. So the angular velocity of a point on the earth is $2\pi/24$ or $\pi/12$ radians per hour. The linear velocity of a point on the surface of the earth depends on its distance from the axis of the earth. What is the linear velocity in miles per hour of a point on the equator? (Use 3950 miles as the radius of the earth).

$$\frac{\pi/12 \text{ rad}}{\text{hr}} \cdot \frac{1 \text{ rev}}{2\pi \text{ rad}} \cdot \frac{(2\pi)(3950) \text{ mi}}{\text{rev}} = \boxed{1034.1 \text{ mph}}$$

The Trigonometric Functions


The six trigonometric functions are the sine (sin), cosine (cos), tangent (tan), cosecant (csc), secant (sec), and cotangent (cot) functions. There are several ways to define these functions of trigonometry. One of the most common mnemonic devices is SOH-CAH-TOA.



$$\sin \alpha = \frac{\text{opposite}}{\text{hypotenuse}} \quad \cos \alpha = \frac{\text{adjacent}}{\text{hypotenuse}} \quad \tan \alpha = \frac{\text{opposite}}{\text{adjacent}}$$

All of these ratios can be written in terms of an angle in the coordinate plane.

If (x, y) is any point other than the origin on the terminal side of an angle α in standard position and $r = \sqrt{x^2 + y^2}$, then




$$\begin{aligned} \sin \alpha &= \frac{\text{opp}}{\text{hyp}} = \frac{y}{r} & \cos \alpha &= \frac{\text{adj}}{\text{hyp}} = \frac{x}{r} & \tan \alpha &= \frac{\text{opp}}{\text{adj}} = \frac{y}{x} \\ \csc \alpha &= \frac{\text{hyp}}{\text{opp}} = \frac{r}{y} & \sec \alpha &= \frac{\text{hyp}}{\text{adj}} = \frac{r}{x} & \cot \alpha &= \frac{\text{adj}}{\text{opp}} = \frac{x}{y} \end{aligned}$$

Note that the cosecant, secant, and cotangent are reciprocals of the sine, cosine, and tangent. These identities are called the reciprocal identities:

$$\csc \alpha = \frac{1}{\sin \alpha} \quad \sec \alpha = \frac{1}{\cos \alpha} \quad \cot \alpha = \frac{1}{\tan \alpha}$$

Examples:


Find the values of the six trigonometric functions of the angle α in standard position whose terminal side passes through $(-2, -4)$.



$$\begin{aligned} r &= \sqrt{2^2 + 4^2} = \sqrt{20} = 2\sqrt{5} \\ x &= -2 \\ y &= -4 \end{aligned}$$

$$\begin{aligned} \sin \alpha &= \frac{y}{r} = \frac{-4}{2\sqrt{5}} = \frac{-2}{\sqrt{5}} & \csc \alpha &= \frac{-\sqrt{5}}{2} \\ \cos \alpha &= \frac{x}{r} = \frac{-2}{2\sqrt{5}} = \frac{-1}{\sqrt{5}} & \sec \alpha &= -\sqrt{5} \\ \tan \alpha &= \frac{y}{x} = \frac{-4}{-2} = 2 & \cot \alpha &= \frac{1}{2} \end{aligned}$$

To find the values of the trigonometric functions of the “special angles” on the unit circle (multiples of 30° and 45°). We could choose any point on the terminal side of each angle and the same ratios would result because the triangles used to calculate the ratios are similar.



$$\sin \alpha = \frac{y}{r} = \frac{y'}{r'} \quad \cos \alpha = \frac{x}{r} = \frac{x'}{r'} \quad \tan \alpha = \frac{y}{x} = \frac{y'}{x'}$$



Since $r = 1$ for any point on the unit circle, points on the unit circle are convenient to use for calculating trigonometry ratios. The question becomes how to find the coordinates of points on the unit circle?

Example: Find the exact values:




- (a) $\sin(90^\circ)$ 1
 (b) $\tan(-5\pi/2)$ undef

- (c) $\sec(180^\circ)$ -1
 (d) $\cot(0)$ undef

We can find the use the ratios that exist in special right triangles to calculate the coordinates of points on the unit circle.



Find the values of x and y in the triangles below:




The *signs* of the trigonometric functions depend on the quadrant in which the angle lies and the corresponding signs of x and y (remember r is always positive).

A good mnemonic to remember which functions are positive in each quadrant is “**All Students Take Calculus**”:


In Quadrant I, all of them are positive, in Quadrant II, \sin (and \csc) are positive, in Quadrant III, \tan (and \cot) are positive, and in Quadrant IV, \cos (and \sec) are positive.

Find the signs of each trigonometric function in each quadrant:



Examples: Find the exact values of the following:

- (1) $\sec 45^\circ = \sqrt{2}$
- (2) $\sin(-3\pi/4) = -\sqrt{2}/2$
- (3) $\tan(5\pi/4) = 1$
- (4) $\cos 135^\circ = -\sqrt{2}/2$
- (5) $\sin(-60^\circ) = -\sqrt{3}/2$
- (6) $\cos(7\pi/6) = -\sqrt{3}/2$
- (7) $\tan 120^\circ = -\sqrt{3}$
- (8) $\csc(-4\pi/3) = 2/\sqrt{3}$
- (9) $\cot(7\pi/3) = 1/\sqrt{3} = \sqrt{3}/3$
- (10) $\sec 210^\circ = -2/\sqrt{3}$



Right Triangle Trigonometry

A solution to the equation $\sin \alpha = \frac{1}{2}$ is an angle whose sine is $\frac{1}{2}$. Because $\sin 30^\circ = \frac{1}{2}$ and $\sin 150^\circ = \frac{1}{2}$, α could be 30° or 150° . Since any angle with the same terminal side as 30° or 150° is also a solution, there are infinitely many solutions. Since right triangles have only acute angles, we are only interested in the acute solutions in this section.

Examples:

Find the angle α that satisfies each equation where $0^\circ \leq \alpha \leq 90^\circ$.

(a) $\sin \alpha = \sqrt{3}/2$

$\alpha = 60^\circ$

(b) $\cos \alpha = 1$

$\alpha = 0^\circ$

(c) $\tan \alpha = 1$

$\alpha = 45^\circ$

Inverse Sine, Cosine, and Tangent Functions

We denote “the angle whose sine is x ” by the symbol $\sin^{-1} x$ (also called the arcsine of x and abbreviated $\arcsin x$) and call it the inverse sine function. Similarly, the inverse cosine and tangent functions are $\cos^{-1} x$ (or $\arccos x$) and $\tan^{-1} x$ (or $\arctan x$) – the angles whose cosine or tangent, respectively, is x .

NOTE: The -1 in $\sin^{-1} x$ does not indicate a reciprocal. $\sin^{-1} x \neq \frac{1}{\sin x}$. The -1 indicates an inverse function. Remember, $\sin^{-1} x$, $\cos^{-1} x$, and $\tan^{-1} x$ are angles!

Because there are infinitely many angles that have a given sine, cosine, or tangent, we define the inverse functions precisely by restricting their domains:

$\sin^{-1} x = \alpha$	provided	$\sin \alpha = x$	and	$-90^\circ \leq \alpha \leq 90^\circ$
$\cos^{-1} x = \alpha$	provided	$\cos \alpha = x$	and	$0^\circ \leq \alpha \leq 180^\circ$
$\tan^{-1} x = \alpha$	provided	$\tan \alpha = x$	and	$-90^\circ < \alpha < 90^\circ$

Examples:

Evaluate each expression. Give the result in degrees. Where necessary, round to the nearest tenth.

(a) $\cos^{-1}(\sqrt{2}/2)$

45°

(b) $\arcsin(\sqrt{3}/2)$


60°

(c) $\tan^{-1}(6.1)$

80.69°

Solving Right Triangles

Finding all the missing angle measures and side lengths of a triangle is called “solving a triangle”. In a right triangle, we usually name the acute angles α and β (beta) and the lengths of the sides opposite those angles a and b , respectively. The 90° angle is γ (gamma) and the length of the side opposite the right angle (the hypotenuse) is c .



To solve a right triangle:

1. Use the Pythagorean Theorem to find the length of a third side when the lengths of two sides are known.
2. Use the trigonometric ratios or inverse trigonometric functions to find missing sides or angles.
3. Use the fact that the sum of the measures of the angles of a triangle is 180° to determine a third angle when two are known.

Examples:

Solve the right triangle in which $\alpha = 60^\circ$ and $c = 2$.




$$\beta = 30^\circ$$

$$\sin 60^\circ = \frac{a}{2} \Rightarrow \frac{\sqrt{3}}{2} = \frac{a}{2} \Rightarrow a = \sqrt{3}$$

$$\begin{aligned} a^2 + b^2 &= c^2 \\ (\sqrt{3})^2 + b^2 &= 2^2 \\ 3 + b^2 &= 4 \\ b^2 &= 1 \end{aligned}$$

$$b = 1$$

Solve the right triangle in which $a = 2$ and $b = 5$.





$$c = \sqrt{2^2 + 5^2} = \sqrt{29}$$


$$\tan \alpha = \frac{2}{5} \Rightarrow \alpha = \tan^{-1}\left(\frac{2}{5}\right) \Rightarrow \alpha = 21.8^\circ$$

$$\begin{aligned} \beta &= 90^\circ - 21.8^\circ \\ \beta &= 68.2^\circ \end{aligned}$$

Using trigonometry, we can find the size of an object without actually measuring the object. Two common terms used in this regard are **angle of elevation** and **angle of depression**.

**Examples:**

The angle of elevation of the top of a cell phone tower is 38.2° at a distance of 344 feet from the tower. What is the height of the tower?




$$\tan 38.2^\circ = \frac{x}{344}$$

$$x = 344 \tan 38.2^\circ$$

$$x = 270.7 \text{ ft}$$

At one location, the angle of elevation of the top of an antenna is 44.2° . At a point that is 100 feet closer to the antenna, the angle of elevation is 63.1° . What is the height of the antenna?



$$\begin{aligned} \tan 63.1^\circ &= \frac{h}{x} & h &= x \tan 63.1^\circ \\ & & h &= 1.971x \end{aligned}$$

$$\tan 44.2^\circ = \frac{h}{x+100}$$

$$h = (\tan 44.2^\circ)(x+100)$$

$$h = (.972)(x+100)$$

$$h = .972x + 97.2$$

$$1.971x = .972x + 97.2$$

$$.999x = 97.2$$

$$x = 97.3$$

$$h = (1.971)(97.3)$$

$$h = 191.8 \text{ ft}$$

The Fundamental Identity and Reference Angles

The Fundamental Identity


The fundamental identity of trigonometry involves the squares of the sine and cosine function. We write $(\cos \alpha)^2$ as $\cos^2 \alpha$ and $(\sin \alpha)^2$ as $\sin^2 \alpha$. Remember that by definition, $\sin \alpha = y/r$, $\cos \alpha = x/r$, and $x^2 + y^2 = r^2$.

$$\sin^2 \alpha + \cos^2 \alpha = \frac{y^2}{r^2} + \frac{x^2}{r^2} = \frac{y^2 + x^2}{r^2} = \frac{r^2}{r^2} = 1$$

The Fundamental Identity of Trigonometry: If α is any angle or real number, then $\sin^2 \alpha + \cos^2 \alpha = 1$.

If we know the sine or cosine of an angle, then we can use the fundamental identity to find the value of the other function of the angle. (Note: you can also figure this out by drawing a triangle and using the Pythagorean Theorem).

Example: Find $\sin \alpha$ given that $\cos \alpha = 1/4$ and α is in Quadrant IV.



$\cos \alpha = \frac{x}{r}$
 $\sin \alpha = \frac{y}{r}$
 $\sin \alpha = \frac{-\sqrt{15}}{4}$

$-OR-$





$\sin^2 \alpha + \cos^2 \alpha = 1$
 $\sin^2 \alpha + (1/4)^2 = 1$
 $\sin^2 \alpha + 1/16 = 1$
 $\sin^2 \alpha = 15/16$
 $\sin \alpha = -\sqrt{15/16}$

Sin is negative in QIV.

 $\sin \alpha = -\frac{\sqrt{15}}{4}$


Reference Angles: When you look at the unit circle, notice that there is a pattern to the coordinates. If you look at all the angles that are 30° away from the x-axis ($30^\circ, 150^\circ, 210^\circ, 330^\circ$), the x-coordinate (cosine) is $\pm\sqrt{3}/2$ and the y-coordinate (sine) is $\pm 1/2$.

Definition: Reference Angle: If θ is a nonquadrantal angle (not on an axis) in standard position, then the reference angle θ' (read “theta prime”) formed by the terminal side of θ and the positive or negative x-axis.




Examples: For each given angle θ , sketch the reference angle θ' and give the measure of θ' in both radians and degrees.


$\theta = 120^\circ$




$\theta = 7\pi/6$



$\theta = 690^\circ$



$\theta = -7\pi/4$



Evaluating Trigonometric Functions Using Reference Angles: For an angle θ in standard position that is not a quadrantal angle:

$$\sin \theta = \pm \sin \theta', \quad \cos \theta = \pm \cos \theta', \quad \tan \theta = \pm \tan \theta',$$

$$\csc \theta = \pm \csc \theta', \quad \sec \theta = \pm \sec \theta', \quad \cot \theta = \pm \cot \theta'$$

where θ' is the reference angle for θ and the sign is determined by the quadrant in which θ lies.

Examples: Find the sine and cosine for each angle using reference angles.

$$\theta = 120^\circ$$

$$\theta = 7\pi/6$$

$$\theta = 690^\circ$$

$$\theta = -7\pi/4$$

$$\theta' = 60^\circ \quad \text{QII}$$

$$\theta' = 30^\circ \quad \text{QIII}$$

$$\theta' = 30^\circ \quad \text{QIV}$$

$$\theta' = \pi/4 \quad \text{QI}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$

$$\boxed{\begin{array}{l} \sin 120^\circ = \frac{\sqrt{3}}{2} \\ \cos 120^\circ = -\frac{1}{2} \end{array}}$$

$$\sin 30^\circ = \frac{1}{2}$$

$$\cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\boxed{\begin{array}{l} \sin 7\pi/6 = -\frac{1}{2} \\ \cos 7\pi/6 = -\frac{\sqrt{3}}{2} \end{array}}$$

$$\sin 30^\circ = \frac{1}{2}$$

$$\cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\boxed{\begin{array}{l} \sin 690^\circ = -\frac{1}{2} \\ \cos 690^\circ = \frac{\sqrt{3}}{2} \end{array}}$$

$$\sin \pi/4 = \frac{\sqrt{2}}{2}$$

$$\cos \pi/4 = \frac{\sqrt{2}}{2}$$

$$\boxed{\begin{array}{l} \sin(-7\pi/4) = \frac{\sqrt{2}}{2} \\ \cos(-7\pi/4) = \frac{\sqrt{2}}{2} \end{array}}$$

Modeling with the Sine Function

The trigonometric functions can be used to model periodic phenomena.

Examples: Demand for a seasonal product can be modeled by the function $d = 200 \sin \frac{\pi(t-3)}{6} + 300$, where d is the number of units sold in month t and where t ranges from 1 through 12. Find the demand in March and June.

$$t=3 \quad t=6$$

$$\text{March: } d = 200 \sin \frac{\pi(3-3)}{6} + 300$$

$$d = 200 \sin 0 + 300$$

$$d = 200(0) + 300$$

$$\boxed{d = 300}$$

$$\text{June: } d = 200 \sin \frac{\pi(6-3)}{6} + 300$$

$$d = 200 \sin \frac{3\pi}{6} + 300$$

$$d = 200 \sin \frac{\pi}{2} + 300$$

$$d = 200(1) + 300$$

$$\boxed{d = 500}$$